

## Systems and Methods for Automated Analysis of Cells and Tissues

1 SVD A11  
Under 35 USC §119(e)(1), this application claims the benefit of prior U.S. provisional application \_\_\_\_\_, filed April 19, 2001, and U.S. provisional application 60/334,723, filed October 31, 2001, the contents of which are incorporated herein by reference. Work described herein was supported in part by funding from the National Institute of Health. The U.S. Government may therefore have certain rights in the invention.

### 1. Background of the Invention

Tissue microarray technology offers the opportunity for high throughput analysis of tissue samples (Konen, J. et al., Nat. Med. 4:844-7 (1998); Kallioniemi, O.P. et al., Hum. Mol. Genet. 10:657-62 (2001); Rimm, D.L. et al., Cancer J. 7:24-31 (2001)). For example, the ability to rapidly perform large scale studies using tissue microarrays can provide critical information for identifying and validating drug targets/ prognostic markers (e.g. estrogen receptor (ER) and HER2/neu) and candidate therapeutics.

Automated quantitative analysis of tissue samples in microarrays, however, presents several challenges, including heterogeneity of tissue sections, subcellular localization of staining, and the presence of background signals. For example, depending on the type of tumor or tissue section being analyzed, the area of interest may represent nearly the entire sample, or only a small percentage. For instance, a pancreatic carcinoma or lobular carcinoma of the breast with substantial desmoplastic response may show stromal tissue representing a large percentage of the total area. If the goal of the assay is to determine epithelial cell expression of a given marker, a protocol must be used that evaluates only that region. The protocol must not only be able to select the region of interest but also normalize it, so that the expression level read from any given area can be compared with that of other areas. Subcellular localization presents similar challenges. Comparisons of nuclear or membranous staining, for example, are quite different from those in total cytoplasmic staining.

Certain methods (including confocal and convolution/ deconvolution microscopy) have been used to quantify expression of proteins at the cellular (or sub-cellular) level within a single high power field (Robinson, J.P. Methods Cell. Biol. 63:89-106 (2001); Shaw, P. Histochem. J. 26:687-94 (1994)). However, these are computationally intensive and laborious techniques, which operate on multiple serial images. As a result, the current standard for analysis of tissue microarrays, like tissue sections, is conventional pathologist-based analysis and grading of the sample according to scale.

Most biomarkers exhibit a parametric (normal, "bell-shaped") distribution, and consequently are best analyzed by a continuous scale (e.g., 0 to 1000). Unfortunately, manual observation tends to be nominal (e.g. 1+, 2+, 3+), primarily because the human eye is unable to reliably distinguish subtle differences in staining intensity. Several methods have been developed to translate nominal manual observations into a continuous scale. Foremost among these is the H-score where the percent of positively stained cells (0 to 100) is multiplied by the staining intensity (e.g. 0 to 3) to make a theoretically continuous scale (0 to 300). However, the inability to detect subtle differences in staining intensity, particularly at the low and high ends of the scale, as well as the tendency to round scores (e.g., 50% at 3+ for a score of 150, versus 47% at 3+ for a score of 141), limits the effectiveness of the H-score.

Automated systems and methods for rapidly analyzing tissue, including tissue microarrays, that permit the identification and localization of identified biomarkers within tissues and other cell containing samples, are needed.

## 2. Summary of the Invention

In one aspect, the invention features systems and methods for rapidly analyzing cell containing samples to localize and quantitate particular biomarkers within cells. In one embodiment, the method is implemented by a computer and superimposes an image of the biomarker against an image of a user defined area within the cell to determine whether the biomarker is within the user defined area.

In another aspect, the invention features an algorithm that facilitates the optical analysis of an array of biological samples, despite image irregularities, distortions, varying topologies, and the absence of one or more elements.

Analysis of patient samples according to the systems and processes described herein can be useful diagnostically (e.g. to identify patients who have a particular disease, have been exposed to a particular toxin or are responding well to a particular therapeutic or organ transplant) and prognostically (e.g. to identify patients who are likely to develop a particular disease, respond well to a particular therapeutic or be accepting of a particular organ transplant). As new and better markers of disease become identified in the post-genomic era, the instant described processes, which not only quantitate the markers, but also determine their relative location within a cell, will increase in applicability.

Automated analysis of cell containing preparations, as described herein, can provide a rapid assessment of the prognostic benefit of biomarkers. In addition, these automated techniques can identify associations that are typically not revealed using manual techniques. Also, automated analysis can better discern subtle differences in staining intensity, particularly at the upper and lower extremes. The ability to detect low level expression and distinguish it from no expression can provide important prognostic information. Furthermore, analysis of the sub-cellular distribution of certain biomarkers may elucidate previously unrecognized associations with patient survival.

Other features, objects, and advantages of the invention will be apparent from the following figures, detailed description and claims.

### 3. Description of the Figures

FIG. 1(A-D) shows separate monochromatic images of a colon carcinoma taken after staining with fluorescently-tagged markers and combined into a single color image as follows: DAPI (to visualize nuclei, blue), anti-cytokeratin (to distinguish tumor from non-tumor elements, green), and anti-alpha-catenin (to visualize cell membranes, red).

FIG. 2(A-D) shows a regression comparison of automated and pathologist-based scoring of estrogen receptor levels.

FIG. 3 is a flowchart of a method for identifying and accounting for the relative location of spots within an array.

FIG. 4 is a flowchart of a process for localizing a signal (e.g. a biomarker) within a locale.

5 FIG. 5 shows a tissue microarray.

FIG. 6 shows an optical microscope station.

#### 4. Detailed Description

##### 4.1 General

10 In general, described herein are a collection of techniques that can be used for rapid, automated analysis of cell containing samples, including tissues and tissue microarrays. While these techniques build on one another and are described as a cohesive process, each technique has wide applicability and may be used individually or in combinations other than those described below.

15 In one embodiment is featured a technique to identify the location of spots within an image. The technique, termed "spotfinder", can flexibly identify such locations despite image irregularities, distortions, varying topologies, and the absence of one or more elements. Although the process is described for locating the position of histospots and identifying missing histospots within tissue microarray images, the technique has broader application. More specifically it can be used to locate elements and identify missing elements in any collection of  
20 elements. Moreover, the process can be used on arrays of virtually any dimension and comprising a variety of elements. The specimens are not limited by size or shape, nor must they be regularly spaced.

25 In another embodiment is featured a technique that can be used alone or in conjunction with spotfinder to optically localize and quantitate a biomarker within a cell. Though an image of a cellular preparation typically features two dimensions, cellular preparations feature depth. For example, one cellular feature may rest atop another. This overlap can potentially confuse image analysis software. A technique described herein, dubbed RESA (Rapid Exponential

Subtraction Algorithm), can approximate a three dimensional image by subtracting out-of-focus image elements. Thus, the impact of background features on an image can be reduced, permitting better image analysis.

Another technique described herein, dubbed PLACE (Pixel Based Locale Assignment  
5 Compartmentalization of Expression), distinguishes between different cellular characteristics. For example, the technique can determine the location of subcellular compartments within individual cells. A computer implementing this technique can, for instance, measure the relative intensities of images derived from compartment-specific stains on a pixel-by-pixel basis. The computer then determines for individual pixels within an image, the likelihood that the pixel  
10 corresponds to a particular locale or user defined area within the cell. Such analysis permits the computer to assign signals to a sub-cellular compartment with an established degree of accuracy (e.g., 95%). The technique can co-localize signals associated with particular biomarkers with images of defined locales within cells.

Use of these techniques can enhance both the speed and accuracy of automated  
15 microarray analysis. Figure 1 shows separate monochromatic images of a colon carcinoma taken after staining with fluorescently-tagged markers and combined into a single color image as follows: DAPI (to visualize nuclei, blue), anti-cytokeratin (to distinguish tumor from non-tumor elements, green), and anti-alpha-catenin (to visualize cell membranes, red) (panel A). Note the significant degree of overlap between the subcellular compartments. A monochromatic image of  
20 the biomarker  $\beta$ -catenin, is taken (panel B, inset) and the intensity of each pixel in the image redistributed according to the relative signal intensity of the various compartments in panel A (blue= nuclei, red= membrane, green = cytoplasm).

Although the  $\beta$ -catenin expression in this tumor is predominantly membrane-associated, the significant overlap in compartments in panel A incorrectly assigns a significant amount of the  
25 signal to the nucleus (magenta and blue pixels, panel B). To aid in the removal of overlapping signals, the monochromatic image of each sub-cellular compartment is exponentially subtracted from an out-of-focus image. Panel C shows a composite of the exponentially subtracted images of DAPI and anti-alpha-catenin (blue and red, respectively), shown on a mask derived from the anti-cytokeratin mask (green pixels). Pixels with too much overlap between channels are  
30 negated (<5%), as are non-tumor areas, as defined by a mask generated from the anti-cytokeratin

image. In panel D, the signal intensity from an exponentially subtracted image of the biomarker ( $\beta$ -catenin, inset) is then redistributed according to the compartments defined in panel C. This results in more accurate assignment of the biomarker to the membrane compartment, which can have important prognostic significance. Since membrane-associated beta-catenin stabilizes cadherin-mediated adhesion by facilitating the cytoskeletal attachment of adhesion complexes, while nuclear-associated beta-catenin acts as a transcription factor, which up-regulates several genes important in cell proliferation and invasion and is considered an oncogene in this capacity, expression of beta-catenin alone does not provide prognostic information. However its localization in the nucleus can be an important indicator of carcinogenesis.

### 1. *Spot-finder*

In one embodiment, shown in FIG. 3, the computer removes B1 any atypically sized spots from the image. Atypically sized spots may include, for example, images of fused spots and/or debris. The computer performs the process automatically, though in other embodiments it may allow use of user input to facilitate the process.

The computer then creates or accesses an opaque virtual mask that is the size and shape of a typical spot. Using the virtual mask, the computer scans B2 the image to determine B3 where the mask first covers an area with the highest average pixel intensity. The computer monitors the total intensity of the image during the scan and, because the mask is opaque, identifies the position of the mask when the total image intensity is minimized. The computer identifies this area as the first spot and sets B4 the pixels within this area to have zero intensity. The computer also sets additional pixels within a predefined area around this area to have zero intensity. This helps to differentiate between overlapping spots.

After identifying the first spot, the computer again scans B2 the image using the mask to find the next area with the highest average pixel intensity. When the next area is found, the computer identifies it as the second spot and sets the pixels in and surrounding this area to have zero intensity. The computer repeats this process until it can no longer find areas of the image with sufficient intensity to qualify as spots.

The computer then identifies B5 a reference point (e.g., the center) in each spot, and draws a line connecting the reference point of each spot to each nearest neighboring spot reference point, above, below, to the left, and to the right. If the computer cannot identify a nearest neighbor in any of these directions (i.e., the spot is on the edge of the array), the computer draws a line from the center of the spot to the nearest edge of the image.

## 2. RESA and PLACE

Once the location of an image area of interest is determined, an optical microscope can obtain a high resolution image at an appropriate wavelength to identify cellular features of interest. These features include the biomarker, also referred to as the “signal”, the cells of interest within the tissue section (referred to as the “cell mask”), or a user defined location within the cell mask, also referred to as the “locale”. The signal, the cell mask, and the locale are referred to as “channels”.

Referring to FIG. 4, a process 50 determines the region of interest in the images by developing a mask from the cell mask channel (step C1). Next, the process applies this mask to the locale and signal channels (step C2). The process then removes out-of-focus information from the masked images, for example, in a process of pseudo-deconvolution (step C3). Next, in a “pixel assignment” phase, the process identifies subcellular features in the image, assigning pixels in the image to the locales (step C4). Once the pixels are assigned, the computer maps the locales onto the signal image (step C5), and quantifies the amount of biomarker in each locale. This phase is referred to as “signal assignment”. These steps are described in greater detail below.

### Masking

During this process, the software identifies a region of interest in the image of the stained cells of interest (i.e., the cell mask channel). The software masks the locale and signal channels avoiding unnecessary analyses of areas outside the region of interest.

To identify a region of interest, the computer determines a threshold intensity for the cell mask channel. Once determined, the computer redistributes the pixel intensities in a binary redistribution. In other words, the computer sets the intensity of each pixel below the threshold

to zero, and sets the remaining pixels to have the maximum intensity (e.g., for an 8-bit image the maximum intensity is 255). The set of pixel locations set to maximum intensity are referred to as the mask. Subsequent procedures on the other images in the image stack are performed on the pixel locations corresponding to the mask.

5 The threshold intensity is related to the intensity of the background in the image, which the computer determines by first binning each pixel according to its intensity (e.g., in an 8-bit image each pixel will have an intensity from 0 to 255). In some embodiments, the background corresponds to the largest bin (i.e., the most common pixel intensity). In other embodiments, the background corresponds to the second largest bin. This occurs in some cases when the tissues autofluoresce and the largest bin corresponds to an area of fluorescing tissue instead of the fluorescing histochemical stains. In either case, the computer assumes that the background intensity is lower than a certain fraction of the maximum intensity (e.g., less than half the maximum intensity).

10 Bin size is plotted versus intensity to yield a histogram. The maximum peak in the histogram corresponds to the largest bin. In embodiments where the largest bin corresponds to the background, the computer assigns the maximum peak intensity as the background intensity. In other embodiments, where the background corresponds to the second largest bin, the histogram has a second peak at a lower intensity than the maximum peak. So, if the second peak is at least a certain fraction of the size of the maximum peak (e.g., at least five percent), then the computer assigns the second peak intensity as the background intensity of the image.

Once established, the computer adds an additional factor to the background intensity to determine the threshold intensity. For an 8-bit image, this factor equals  $D(1/5)/10$  multiplied by a user defined input (usually 0.5). Here,  $D(1/5)$  is the quintile distribution of the binned pixels, which is determined as

$$25 \quad D(1/5) = \langle I \rangle_{top20} - \langle I \rangle_{bottom20},$$

where  $\langle I \rangle_{top20}$  is the mean pixel intensity of the pixels within the top 20<sup>th</sup> percentile, and  $\langle I \rangle_{bottom20}$  is the mean pixel intensity of pixels in the bottom 20<sup>th</sup> percentile.

Pixels with intensity at or above the threshold intensity are assigned to the mask.

The mask is then further modified according to user-defined parameters and image processing techniques. For example, the mask can be dilated or eroded so that the mask area matches a user-defined value, or have holes of a particular (user-defined) size within it filled.

5 The user-defined parameters for creating the mask may be adjusted after analyzing a small number of sample histospot images, prior to running the entire array.

After developing the mask, the computer applies the mask to the images in the image stack, identifying the region of interest in each of these images as the pixel locations corresponding to the mask pixel locations.

### 10 Background Reduction

As shown, the process 50 reduces C3 the impact of the out-of-focus information from the image. For example, the process 50 may use a pseudo deconvolution technique. While the pixels of the remaining image are reduced in intensity, the image information represents a thinner virtual slice through the top of the tissue. Furthermore, pseudo-deconvolution enhances the interfacial areas between the higher stain intensity and lower stain intensity areas of the image by increasing the contrast between these areas.

20 The computer performs pseudo-deconvolution on the locale (cellular compartments) and signal (i.e., cellular components) channels. The computer first masks the images of these channels, reducing the number of pixels to be analyzed. The computer analyzes two images of each channel. The first image is an in-focus image (i.e., an image of the top of the histospot). The second image is a slightly out-of-focus image, produced by placing the focal plane slightly below the bottom of the tissue (e.g., for a five micron thick histospot, the focal plane of this image is located about eight microns below the top of the histospot).

25 For each pixel location, the computer subtracts a percentage of the out-of-focus image pixel intensity,  $I_{out-of-focus}$ , from the corresponding in-focus image pixel intensity,  $I_{in-focus}$ . The computer determines the adjusted pixel intensity,  $I_{new\ pixel}$ , using the quartile distribution,  $D(\frac{1}{4})$ , of the in-focus image as follows:

$$I_{\text{new pixel}} = I_{\text{in-focus}} - I_{\text{out-of-focus}} \times \left( \frac{I_{\text{max}} - I_{\text{in-focus}}}{I_{\text{max}}} \right)^{\psi},$$

where  $I_{\text{max}}$  is the maximum pixel intensity (e.g., 255 for an 8-bit image), and  $\psi$  is calculated from

$$\psi = \alpha \times D(\mathcal{V}_4)^{-\beta},$$

which was developed from an empiric assessment of a library of images. Optical deconvolutions were judged visually and the  $\psi$  for each was plotted versus the quartile distribution for the in-focus image. Regression analysis of the empiric data yielded values for the fitting-parameters (i.e.,  $\alpha$  is about 80 and  $\beta$  is about 1.19). The quartile distribution is determined from

$$D(\mathcal{V}_4) = \langle I \rangle_{\text{top}25} - \langle I \rangle_{\text{bottom}25},$$

where  $\langle I \rangle_{\text{top}25}$  is the mean pixel intensity of the pixels within the top 25<sup>th</sup> percentile, and  $\langle I \rangle_{\text{bottom}25}$  is the mean pixel intensity of pixels in the bottom 25<sup>th</sup> percentile. Conceptually, low intensity pixels in images with a low  $D(\mathcal{V}_4)$  (i.e. a low signal to noise ratio) are subtracted less heavily than low intensity pixels from images with a high  $D(\mathcal{V}_4)$ .

The value of  $\psi$  may be refined by determining the percent of signal intensity remaining after pseudo-deconvolution within the masked area and comparing it to a predefined value for that channel. If the percent is, for example, greater than the predefined value then the pseudo-deconvolution stops. Otherwise, the computer iteratively increases the value of  $\psi$  until the predefined percent of signal intensity is reached. The predefined value is the expected percentage of the mask covered by a channel.

After pseudo-deconvolution, each pixel of the resulting images is assigned to a locale in a process referred to as pixel assignment.

#### Pixel Based Locale Assignment Compartmentalization of Expression (PLACE)

During the pixel assignment phase, the computer assigns an identity based on the relative intensity of that pixel location in each of the locale channel images (i.e., the images of the stained

locales). For example, during this phase the computer decides for each pixel location in the image whether it belongs to the nucleus, the membrane, or the cytoplasm. The computer does not make an assignment to pixels that it cannot assign within a user-defined degree of confidence (e.g., 95%). Higher levels of confidence eliminate more pixels from the analysis.

5 In general, for each pixel location in two locale images the computer reads a pixel intensity and compares each intensity value to a predetermined threshold intensity value. If the intensity value of only one locale is greater than the threshold, the computer assigns the pixel location to that locale. If both the intensity values are greater than their respective thresholds, the computer compares the intensity values from each locale, and assigns the identity of the locale having the greater intensity to that pixel location. If both the pixel intensities are below their threshold values, the computer assigns the pixel to a third locale.

10 After repeating the above for pixel locations in the images, the computer calculates the area of each locale, and compares the result to a predetermined (expected) coverage fraction. If the calculated coverage fraction (e.g., number of nuclear locale pixels/number of masked pixels) is greater than the predetermined coverage fraction, then the computer removes the pixels having the lowest intensity from the locale. The computer continues to remove the lowest intensity pixels until the coverage fraction is reduced to about the predetermined coverage fraction.

15 The following is an example of how this process works. The membrane locale and the nucleus locale images are selected for assignment analysis performed at 95% confidence interval. Pixel locations are assigned to the cytoplasm locale by exclusion.

20 The computer reads pixel intensities at each pixel location in the membrane and nucleus locale images, and compares them to threshold values. If neither of the intensity values are greater than the threshold values, the pixel location is assigned to the cytoplasm locale. If only the nuclear intensity or membrane intensity is greater than the threshold value, the computer assigns the pixel location to the above-threshold locale. If both intensities are higher than the thresholds, computer compares the ratio of the intensity values to one, and makes an assignment as follows:

$$\frac{\text{nuclear intensity}}{\text{membrane intensity}} > 1 \rightarrow \text{pixel location} = \text{nuclear locale}$$

$$\frac{\text{nuclear intensity}}{\text{membrane intensity}} < 1 \rightarrow \text{pixel location} = \text{membrane locale}.$$

$$\frac{\text{nuclear intensity}}{\text{membrane intensity}} = 1 \rightarrow \text{pixel location} = \text{unassigned}$$

Thus, if the nuclear intensity is greater than the membrane intensity, the computer assigns the pixel location to the nuclear locale. If the membrane intensity is greater than the nuclear intensity, the computer assigns the pixel location to the membrane locale. If the membrane intensity is equal to the nuclear intensity, the pixel location is unassigned. This repeats for the pixel locations.

Once all the pixel locations have been analyzed, the computer determines the amount of nuclear intensity incorrectly assigned to the membrane locale (i.e., nuclear to membrane spillover), and vice versa. If the amount of nuclear intensity incorrectly assigned to the membrane channel is >5% of the total nuclear intensity, then the computer weights the nuclear intensity by a factor,  $w$ , and recalculates the ratio of weighted nuclear intensity to membrane intensity. This ratio is compared to one, and pixel locations are reassigned as follows:

$$\frac{w \times \text{nuclear intensity}}{\text{membrane intensity}} > 1 \rightarrow \text{pixel location} = \text{nuclear locale}$$

$$\frac{w \times \text{nuclear intensity}}{\text{membrane intensity}} < 1 \rightarrow \text{pixel location} = \text{membrane locale}.$$

$$\frac{w \times \text{nuclear intensity}}{\text{membrane intensity}} = 1 \rightarrow \text{pixel location} = \text{unassigned}$$

The computer again determines the amount of each locale incorrectly assigned. If this is still >5% the computer increases the value of  $w$  and reiterates the steps above. This continues until the amount of incorrectly assigned nuclear locale is <5%. The computer employs a similar technique to minimize the membrane-to-nuclear spillover.

The computer also calculates the area of the cytoplasmic (exclusion) locale and compares it to a predetermined value. By iterating the assignment process, the computer ensures that there is <5% cytoplasmic-to-nuclear or cytoplasmic-to-membrane, as determined based on the biology.

The computer then evaluates the amount of signal in each locale during a “signal assignment” process.

Following pixel assignment, the computer sums the signal in each locale. The computer reads the pixel intensity of the signal image (i.e., the image of the stain that selectively labels the cellular component), and adds together the signal intensity for pixel locations assigned to like subcellular compartments. The computer calculates a pixel intensity sum of a locale by the direct addition of the signal intensity of each pixel location assigned to that locale. The computer also calculates a sum of pixel intensity ratios by adding together the ratio of the signal intensity and the locale intensity for each pixel location.

The pixel intensity sum and pixel intensity ratio sum is then used in calculating one or more parameters. For example, the computer determines the relative percentage of signal falling within each of the compartments (e.g. 30% of the total signal is membranous, 20% is cytoplasmic, and 50% is nuclear). In another example, the computer expresses the amount of signal present relative to the size of a particular compartment (e.g. the signal intensity of pixels assigned to the membrane channel divided by the number of pixels assigned to the membrane channel). The user may select to have the computer evaluate other parameters of interest. For example, how much of the image area is covered by the mask, how much of the mask is covered by each locale, etc.

By implementing the pseudo-deconvolution algorithm (which limits the majority of extraneous pixel intensity) together with intensity area measurements (which further define the area of a particular sub-cellular locale), the computer is able to make highly accurate assignments of pixel locations sub-cellular locations.

In some embodiments, the computer performs additional steps to better utilize the dynamic range of the camera. This is achieved by redistributing the pixel intensities in an image across the dynamic range of the detector based on their relative intensities.

One form of redistribution is normalized redistribution, whereby the lower threshold (i.e., the pixel intensity of the background, determined during masking) is subtracted from all the pixels in the image, and any pixel with a resulting negative value is set to zero. Normalized

redistribution is used for the signal channel as this redistribution preserves the scale from one sample to the next, allowing direct comparisons to be made between samples. This is performed either after masking the signal image.

Double-logarithmic redistribution sets all pixels in an image above 50% of the image's upper threshold (i.e., the value which only 50% of the pixels in the image have greater intensity) to the maximum intensity value (e.g., 255 for an 8-bit image). All pixels with intensity values below the lower threshold are set to 0, and all pixels with intensity values between the upper and lower thresholds are reassigned according to the formula:

$$I_{\text{new}} = \frac{\log(I_{\text{old}} - LT)}{I_{\text{max}} \log(\frac{1}{2}UT - LT)},$$

where  $I_{\text{new}}$  refers to the new pixel intensity,  $I_{\text{old}}$  refers to the old pixel intensity,  $LT$  and  $UT$  are the lower and 50% maximum thresholds, respectively, and  $I_{\text{max}}$  is the maximum intensity value.

Double-logarithmic redistribution is used for the locale channels, either after masking or after pseudo-deconvolution of these channels. Conceptually, it ensures that pixels in locale images that have intensities above the 50<sup>th</sup> percentile are assigned to their locale during the assignment phase. Pixels with intensities below, but close to, the 50<sup>th</sup> percentile are weighted more heavily and are more likely to be assigned to the locale than pixels that have intensities well below the 50<sup>th</sup> percentile.

Other user-defined redistributions, such as linear redistributions or other equation-based redistributions, may be used in addition to the above-described examples.

Although the algorithms described above are with reference to analysis of tissue microarrays, they are not limited to studying only such arrays. The spotfinder algorithm may be used for identifying the location of any element comprising a collection and the RESA and PLACE algorithms may be used to localize and quantitate a biomarker within any imageable, cell containing sample, including tissue biopsies and cell containing fluid samples, such as blood, urine, spinal fluid, saliva, lymph, pleural fluid, peritoneal fluid and pericardial fluid.

Also, any optical or non-optical imaging device can be used, such as for example, upright or inverted optical microscopes, scanning confocal microscopes, cameras, scanning or tunneling electron microscopes, scanning probe microscopes, and imaging infrared detectors etc.

5 In the embodiments described above, the computer can include hardware, software, or a combination of both to control the other components of the system and to analyze the images to extract the desired information about the histospots and tissue microarrays. The analysis described above is implemented in computer programs using standard programming techniques. Such programs are designed to execute on programmable computers each comprising a processor, a data storage system (including memory and/or storage elements), at least one input device, at least one output device, such as a display or printer. The program code is applied to input data (e.g., stitched together images or image stacks) to perform the functions described herein and generate information (e.g., localization of signal), which is applied to one or more output devices. Each computer program can be implemented in a high-level procedural or object-oriented programming language, or an assembly or machine language. Each such computer program can be stored on a computer readable storage medium (e.g., CD ROM or magnetic diskette) that when read by a computer can cause the processor in the computer to perform the analysis described herein.

10 The following provides a detailed description of a specific embodiment of the preparation and analysis of tissue microarrays according to methods described herein, although similar steps could be performed with respect to any cell containing sample. Referring to FIG. 5, a tissue microarray 100 includes multiple samples of histospots 120 prepared from histocores embedded typically in a thin (e.g., about five microns) block of paraffin 130 at regular intervals, forming a series of rows and columns. Histospots (thin sections of histocores) 120 may be substantially disk-like in shape and will typically have the same thickness as the paraffin block 130 (i.e., about 15 five microns) and a diameter of about 0.6 millimeters. Typically the centers of the histospots are spaced about a few tenths of a millimeter apart. Paraffin block 130 and histospots 120 may be mounted on a microscope slide 110. A tissue microarray 100 may include any number of histospots, typically on the order of several hundred to a few thousand.

20 Referring to FIG. 6, an optical microscopy station can be used to obtain an appropriate image of the tissue. Microscopy station 200 includes an inverted optical microscope 201 for

imaging the tissue, and a computer 290 for analyzing the images. Optical microscope 201 includes a mount 210, housing a light source 220, a sample stage 240, an objective lens 250 and a CCD camera 270. A frame grabber in computer 290 acquires the images through CCD camera 270.

5           Optical microscope 201 also includes filter wheels 230 and 260, which house a series of dichroic filters. The filters in wheel 230 allow selection of the appropriate illumination spectra for standard or fluorescent microscopy. Filters in wheel 260 filter the transmitted light for isolation of spectral signatures in fluorescent microscopy. Sample stage 240 supports and appropriately positions tissue microarray 100. Sample stage 240 can be linearly translated in the x, y, and z directions (axes are shown). Sample stage 240 includes motors to enable automated translation. Computer 290 controls sample stage 240 translation by servo control of the motors.

10           A tissue microarray 100 can be imaged as follows: a user places the microarray on a sample stage 240. The user adjusts sample stage 240 so that the first (i.e., top-left) histospot is at the center of the field of view and focused on CCD camera 270. The objective lens 250 should be adjusted to the appropriate resolution, for example, a 0.6 millimeter histospot can be viewed at 10X magnification. Generally, the histospots correspond to areas of higher light intensity than the surrounding paraffin, as assessed through various means including signals derived from the visible light scattering of stained tissues, tissue autofluorescence or from a fluorescent tag.

15           Computer 290 can acquire a low-resolution image (e.g. 64 pixel x 64 pixel with 16 bin resolution) using computer software (Softworx 2.5, Applied Precision, Issaquah, WA) and an imaging platform (e.g., Deltavision). Computer 290 automatically translates sample stage 240 by an amount approximately equal to a field of view. The computer then acquires a second low-resolution image. This process is repeated until the computer has acquired images of the entire tissue microarray. Then, using commercially available software, the computer generates a

20           composite image of the entire tissue microarray by stitching together the sequence of images like a patchwork.

25

30           Biological markers, which may be detected in accordance with the present invention include, but are not limited to any nucleic acids, proteins, peptides, lipids, carbohydrates or other components of a cell. Certain markers are characteristic of particular cells, while other markers have been identified as being associated with a particular disease or condition. Examples of

known prognostic markers include enzymatic markers such as galactosyl transferase II, neuron specific enolase, proton ATPase-2, and acid phosphatase. Hormone or hormone receptor markers include human chorionic gonadotropin (HCG), adrenocorticotrophic hormone, carcinoembryonic antigen (CEA), prostate-specific antigen (PSA), estrogen receptor, progesterone receptor, androgen receptor, gC1q-R/p33 complement receptor, IL-2 receptor, p75 neurotrophin receptor, PTH receptor, thyroid hormone receptor, and insulin receptor.

Lymphoid markers include alpha-1-antichymotrypsin, alpha-1-antitrypsin, B cell marker, bcl-2, bcl-6, B lymphocyte antigen 36kD, BM1 (myeloid marker), BM2 (myeloid marker), galectin-3, granzyme B, HLA class I Antigen, HLA class II (DP) antigen, HLA class II (DQ) antigen, HLA class II (DR) antigen, human neutrophil defensins, immunoglobulin A, immunoglobulin D, immunoglobulin G, immunoglobulin M, kappa light chain, kappa light chain, lambda light chain, lymphocyte/histocyte antigen, macrophage marker, muramidase (lysozyme), p80 anaplastic lymphoma kinase, plasma cell marker, secretory leukocyte protease inhibitor, T cell antigen receptor (JOVI 1), T cell antigen receptor (JOVI 3), terminal deoxynucleotidyl transferase, unclustered B cell marker.

Tumour markers include alpha fetoprotein, apolipoprotein D, BAG-1 (RAP46 protein), CA19-9 (sialyl lewisa), CA50 (carcinoma associated mucin antigen), CA125 (ovarian cancer antigen), CA242 (tumour associated mucin antigen), chromogranin A, clusterin (apolipoprotein J), epithelial membrane antigen, epithelial-related antigen, epithelial specific antigen, gross cystic disease fluid protein-15, hepatocyte specific antigen, heregulin, human gastric mucin, human milk fat globule, MAGE-1, matrix metalloproteinases, melan A, melanoma marker (HMB45), mesothelin, metallothionein, microphthalmia transcription factor (MITF), Muc-1 core glycoprotein, Muc-1 glycoprotein, Muc-2 glycoprotein, Muc-5AC glycoprotein, Muc-6 glycoprotein, myeloperoxidase, Myf-3 (Rhabdomyosarcoma marker), Myf-4 (Rhabdomyosarcoma marker), MyoD1 (Rhabdomyosarcoma marker), myoglobin, nm23 protein, placental alkaline phosphatase, prealbumin, prostate specific antigen, prostatic acid phosphatase, prostatic inhibin peptide, PTEN, renal cell carcinoma marker, small intestinal mucinous antigen, tetranectin, thyroid transcription factor-1, tissue inhibitor of matrix metalloproteinase 1, tissue inhibitor of matrix metalloproteinase 2, tyrosinase, tyrosinase-related protein-1, villin, von Willebrand factor.

Cell cycle associated markers include apoptosis protease activating factor-1, bcl-w , bcl-x, bromodeoxyuridine, CAK (cdk-activating kinase), cellular apoptosis susceptibility protein (CAS), caspase 2, caspase 8, CPP32 (caspase-3), cyclin dependent kinases, cyclin A, cyclin B1, cyclin D1, cyclin D2, cyclin D3, cyclin E, cyclin G, DNA fragmentation factor (N-terminus), Fas (CD95), Fas-associated death domain protein, Fas ligand, Fen-1, IPO-38, Mcl-1, minichromosome maintenance proteins, mismatch repair protein (MSH2), poly (ADP-Ribose) polymerase, proliferating cell nuclear antigen, p16 protein, p27 protein, p34cdc2, p57 protein (Kip2), p105 protein, Stat 1 alpha, topoisomerase I, topoisomerase II alpha, topoisomerase III alpha, topoisomerase II beta.

Neural tissue and tumour markers include alpha B crystallin, alpha-internexin, alpha synuclein, amyloid precursor protein, beta amyloid, calbindin, choline acetyltransferase, excitatory amino acid transporter 1, GAP43, glial fibrillary acidic protein, glutamate receptor 2, myelin basic protein, nerve growth factor receptor (gp75), neuroblastoma marker, neurofilament 68kD, neurofilament 160kD, neurofilament 200kD, neuron specific enolase, nicotinic acetylcholine receptor alpha4, nicotinic acetylcholine receptor beta2, peripherin, protein gene product 9, S-100 protein, serotonin, SNAP-25, synapsin I, synaptophysin, tau, tryptophan hydroxylase, tyrosine hydroxylase, ubiquitin.

Cluster differentiation markers include CD1a, CD1b, CD1c, CD1d, CD1e, CD2, CD3delta, CD3epsilon, CD3gamma, CD4, CD5, CD6, CD7, CD8alpha, CD8beta, CD9, CD10, CD11a, CD11b, CD11c, CDw12, CD13, CD14, CD15, CD15s, CD16a, CD16b, CDw17, CD18, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD26, CD27, CD28, CD29, CD30, CD31, CD32, CD33, CD34, CD35, CD36, CD37, CD38, CD39, CD40, CD41, CD42a, CD42b, CD42c, CD42d, CD43, CD44, CD44R, CD45, CD46, CD47, CD48, CD49a, CD49b, CD49c, CD49d, CD49e, CD49f, CD50, CD51, CD52, CD53, CD54, CD55, CD56, CD57, CD58, CD59, CDw60, CD61, CD62E, CD62L, CD62P, CD63, CD64, CD65, CD65s, CD66a, CD66b, CD66c, CD66d, CD66e, CD66f, CD68, CD69, CD70, CD71, CD72, CD73, CD74, CDw75, CDw76, CD77, CD79a, CD79b, CD80, CD81, CD82, CD83, CD84, CD85, CD86, CD87, CD88, CD89, CD90, CD91, CDw92, CDw93, CD94, CD95, CD96, CD97, CD98, CD99, CD100, CD101, CD102, CD103, CD104, CD105, CD106, CD107a, CD107b, CDw108, CD109, CD114, CD115, CD116, CD117, CDw119, CD120a, CD120b, CD121a, CDw121b, CD122, CD123, CD124, CDw125,

CD126, CD127, CDw128a, CDw128b, CD130, CDw131, CD132, CD134, CD135, CDw136, CDw137, CD138, CD139, CD140a, CD140b, CD141, CD142, CD143, CD144, CDw145, CD146, CD147, CD148. CDw149, CDw150, CD151, CD152, CD153, CD154, CD155, CD156, CD157, CD158a, CD158b, CD161, CD162, CD163, CD164, CD165, CD166, and TCR-zeta.

5           Other cellular markers include centromere protein-F (CENP-F), giantin, involucrin, lamin A&C [XB10], LAP-70, mucin, nuclear pore complex proteins, p180 lamellar body protein, ran, r, cathepsin D, Ps2 protein, Her2-neu, P53, S100, epithelial marker antigen (EMA), TdT, MB2, MB3, PCNA, and Ki67.

10           Cell containing samples may be stained using dyes or stains, or histochemicals, that directly react with the specific biomarkers or with various types of cells or subcellular compartments. Not all stains are compatible. Therefore the type of stains employed and their sequence of application should be well considered, but can be readily determined by one of skill in the art. Such histochemicals may be chromophores detectable by transmittance microscopy or fluorophores detectable by fluorescence microscopy. In general, a cell containing samples may be incubated with a solution comprising at least one histochemical, which will directly react with or bind to chemical groups of the target. Some histochemicals must be co-incubated with a mordant, or metal, in order to allow staining. A cell containing sample may be incubated with a mixture of at least one histochemical that stains a component of interest and another histochemical that acts as a counterstain and binds a region outside the component of interest.

15           Alternatively, mixtures of multiple probes may be used in the staining, and provide a way to identify the positions of specific probes.

20

          The following, non-limiting list provides exemplary chromophores that may be used as histological stains or counterstains and their target cells, subcellular compartments, or cellular components: Eosin (alkaline cellular components, cytoplasm), Hematoxylin (nucleic acids),

25           Orange G (red blood, pancreas, and pituitary cells), Light Green SF (collagen), Romanowsky-Giemsa (overall cell morphology), May-Grunwald (blood cells), Blue Counterstain (Trevigen), Ethyl Green (CAS) (amyloid), Feulgen-Naphthol Yellow S (DNA), Giemsa (differentially stains various cellular compartments), Methyl Green (amyloid), pyronin (nucleic acids), Naphthol-

30           Yellow (red blood cells), Neutral Red (nuclei), Papanicolaou stain (which typically includes a

mixture of Hematoxylin, Eosin Y, Orange G and Bismarck Brown mixture (overall cell morphology), Red Counterstain B (Trevigen), Red Counterstain C (Trevigen), Sirius Red (amyloid), Feulgen reagent (pararosanilin) (DNA), Gallocyanin chrom-alum (DNA), Gallocyanin chrom-alum and Naphthol Yellow S (DNA), Methyl Green-Pyronin Y (DNA), Thionin-Feulgen reagent (DNA), Acridine Orange (DNA), Methylene Blue (RNA and DNA), Toluidine Blue (RNA and DNA), Alcian blue (carbohydrates), Ruthenium Red (carbohydrates), Sudan Black (lipids), Sudan IV (lipids), Oil Red-O (lipids), Van Gieson's trichrome stain (acid fuchsin and picric acid mixture ) (muscle cells), Masson trichrome stain (hematoxylin, acid fuchsin, and Light Green mixture) (stains collagen, cytoplasm, nucleoli differently), Aldehyde Fuchsin (elastin fibers), and Weigert stain (differentiates reticular and collagenous fibers). A comprehensive list of such stains, their description, and general use is given in R.D. Lillie, "Conn's Biological Stains", 8th ed., Williams and Wilkins Company, Baltimore, Maryland (1969). Suitable mordants and compositions of the preceding are well-known to one of skill in the art.

The following, non-limiting list provides exemplary fluorescent histological stains and their target cells, subcellular compartments, or cellular components if applicable: 4',6-diamidino-2-phenylindole (DAPI) (nucleic acids), Eosin (alkaline cellular components, cytoplasm), Hoechst 33258 and Hoechst 33342 (two bisbenzimidides) (nucleic acids), Propidium Iodide (nucleic acids), Spectrum Orange (nucleic acids), Spectrum Green (nucleic acids), Quinacrine (nucleic acids), Fluorescein-phalloidin (actin fibers), Chromomycin A 3 (nucleic acids), Acriflavine-Feulgen reaction (nucleic acid), Auramine O-Feulgen reaction (nucleic acids), Ethidium Bromide (nucleic acids). Nissl stains (neurons), high affinity DNA fluorophores such as POPO, BOBO, YOYO and TOTO and others, and Green Fluorescent Protein fused to DNA binding protein, such as histones, ACMA, Quinacrine and Acridine Orange.

A wide variety of proprietary fluorescent organelle-specific probes are available from Molecular Probes (Eugene, OR), which include mitochondria-specific probes (MitoFluor and MitoTracker dyes), endoplasmic reticulum (ER) and Golgi probes (ER-Tracker and various ceramide conjugates), and lysosomal probes (LysoTracker dyes). These probes, as well as many nonproprietary fluorescent histochemicals, are available from and extensively described in the

*Handbook of Fluorescent Probes and Research Products* 8<sup>th</sup> Ed. (2001), available from Molecular Probes, Eugene, OR.

Each cell containing sample may be co-incubated with appropriate substrates for an enzyme that is a cellular component of interest and appropriate reagents that yield colored precipitates at the sites of enzyme activity. Such enzyme histochemical stains are specific for the particular target enzyme. Staining with enzyme histochemical stains may be used to define a subcellular component or a particular type of cell. Alternatively, enzyme histochemical stains may be used diagnostically to quantitate the amount of enzyme activity in cells. A wide variety of enzymatic substrates and detection assays are known and described in the art, and some selected methods are exemplified below.

Acid phosphatases may be detected through several methods. In the Gomori method for acid phosphatase, a cell preparation is incubated with glycerophosphate and lead nitrate. The enzyme liberates phosphate, which combines with lead to produce lead phosphate, a colorless precipitate. The tissue is then immersed in a solution of ammonium sulfide, which reacts with lead phosphate to form lead sulfide, a black precipitate. Alternatively, cells may be incubated with a solution comprising pararosanilin-HCl, sodium nitrite, naphthol ASB1 phosphate (substrate), and veronal acetate buffer. This method produces a red precipitate in the areas of acid phosphatase activity. Owing to their characteristic content of acid phosphatase, lysosomes can be distinguished from other cytoplasmic granules and organelles through the use of this assay.

Dehydrogenases may be localized by incubating cells with an appropriate substrate for the species of dehydrogenase and tetrazole. The enzyme transfers hydrogen ions from the substrate to tetrazole, reducing tetrazole to formazan, a dark precipitate. For example, NADH dehydrogenase is a component of complex I of the respiratory chain and is localized predominantly to the mitochondria.

Other enzymes for which well-known staining techniques have been developed, and their primary cellular locations or activities, include but are not limited to the following: ATPases

(muscle fibers), succinate dehydrogenases (mitochondria), cytochrome c oxidases (mitochondria), phosphorylases (mitochondria), phosphofructokinases (mitochondria), acetyl cholinesterases (nerve cells), lactases (small intestine), leucine aminopeptidases (liver cells), myodenylate deaminases (muscle cells), NADH diaphorases (erythrocytes), and sucrases (small intestine).

Immunohistochemistry is among the most sensitive and specific histochemical techniques. Each histospot may be combined with a labeled binding composition comprising a specifically binding probe. Various labels may be employed, such as fluorophores, or enzymes which produce a product which absorbs light or fluoresces. A wide variety of labels are known which provide for strong signals in relation to a single binding event. Multiple probes used in the staining may be labeled with more than one distinguishable fluorescent label. These color differences provide a way to identify the positions of specific probes. The method of preparing conjugates of fluorophores and proteins, such as antibodies, is extensively described in the literature and does not require exemplification here.

Although there are at least 120,000 commercially available antibodies, the following lists some exemplary primary antibodies known to specifically bind cellular components and which are presently employed as components in immunohistochemical stains used for research and, in limited cases, for diagnosis of various diseases. Anti-estrogen receptor antibody (breast cancer), anti-progesterone receptor antibody (breast cancer), anti-p53 antibody (multiple cancers), anti-Her-2/neu antibody (multiple cancers), anti-EGFR antibody (epidermal growth factor, multiple cancers), anti-cathepsin D antibody (breast and other cancers), anti-Bcl-2 antibody (apoptotic cells), anti-E-cadherin antibody, anti-CA125 antibody (ovarian and other cancers), anti-CA15-3 antibody (breast cancer), anti-CA19-9 antibody (colon cancer), anti-c-erbB-2 antibody, anti-P-glycoprotein antibody (MDR, multi-drug resistance), anti-CEA antibody (carcinoembryonic antigen), anti-retinoblastoma protein (Rb) antibody, anti-ras oncoprotein (p21) antibody, anti-Lewis X (also called CD15) antibody, anti-Ki-67 antibody (cellular proliferation), anti-PCNA (multiple cancers) antibody, anti-CD3 antibody (T-cells), anti-CD4 antibody (helper T cells), anti-CD5 antibody (T cells), anti-CD7 antibody (thymocytes, immature T cells, NK killer cells), anti-CD8 antibody (suppressor T cells), anti-CD9/p24 antibody (ALL), anti-CD10 (also called

CALLA) antibody (common acute lymphoblastic leukemia), anti-CD11c antibody (Monocytes, granulocytes, AML), anti-CD13 antibody (myelomonocytic cells, AML), anti-CD14 antibody (mature monocytes, granulocytes), anti-CD15 antibody (Hodgkin's disease), anti-CD19 antibody (B cells), anti-CD20 antibody (B cells), anti-CD22 antibody (B cells), anti-CD23 antibody (activated B cells, CLL), anti-CD30 antibody (activated T and B cells, Hodgkin's disease), anti-CD31 antibody (angiogenesis marker), anti-CD33 antibody (myeloid cells, AML), anti-CD34 antibody (endothelial stem cells, stromal tumors), anti-CD35 antibody (dendritic cells), anti-CD38 antibody (plasma cells, activated T, B, and myeloid cells), anti-CD41 antibody (platelets, megakaryocytes), anti-LCA/CD45 antibody (leukocyte common antigen), anti-CD45RO antibody (helper, inducer T cells), anti-CD45RA antibody (B cells), anti-CD39, CD100 antibody, anti-CD95/Fas antibody (apoptosis), anti-CD99 antibody (Ewings Sarcoma marker, MIC2 gene product), anti-CD106 antibody (VCAM-1; activated endothelial cells), anti-ubiquitin antibody (Alzheimer's disease), anti-CD71 (transferrin receptor) antibody, anti-c-myc (oncoprotein and a hapten) antibody, anti-cytokeratins (transferrin receptor) antibody, anti-vimentins (endothelial cells) antibody (B and T cells), anti-HPV proteins (human papillomavirus) antibody, anti-kappa light chains antibody (B cell), anti-lambda light chains antibody (B cell), anti-melanosomes (HMB45) antibody (melanoma), anti-prostate specific antigen (PSA) antibody (prostate cancer), anti-S-100 antibody (melanoma, salivary, glial cells), anti-tau antigen antibody (Alzheimer's disease), anti-fibrin antibody (epithelial cells), anti-keratins antibody, anti-cytokeratin antibody (tumor), anti-alpha-catenin (cell membrane), and anti-Tn-antigen antibody (colon carcinoma, adenocarcinomas, and pancreatic cancer).

Fluorophores that may be conjugated to a primary antibody include but are not limited to Fluorescein, Rhodamine, Texas Red, Cy2, Cy3, Cy5, VECTOR Red, ELF.TM. (Enzyme-Labeled Fluorescence), Cy0, Cy0.5, Cy1, Cy1.5, Cy3, Cy3.5, Cy5, Cy7, FluorX, Calcein, Calcein-AM, CRYPTOFLUOR.TM.'S, Orange (42 kDa), Tangerine (35 kDa), Gold (31 kDa), Red (42 kDa), Crimson (40 kDa), BHMP, BHDMAP, Br-Oregon, Lucifer Yellow, Alexa dye family, N-[6-(7-nitrobenz-2-oxa-1, 3-diazol-4-yl)amino]caproyl] (NBD), BODIPY.TM., boron dipyrromethene difluoride, Oregon Green, MITOTRACKER.TM. Red, DiOC.sub.7 (3), DiIC.sub.18, Phycoerythrin, Phycobiliproteins BPE (240 kDa) RPE (240 kDa) CPC (264 kDa) APC (104 kDa), Spectrum Blue, Spectrum Aqua, Spectrum Green, Spectrum Gold, Spectrum Orange,

Spectrum Red, NADH, NADPH, FAD, Infra-Red (IR) Dyes, Cyclic GDP-Ribose (cGDPR), Calcofluor White, Lissamine, Umbelliferone, Tyrosine and Tryptophan. A wide variety of other fluorescent probes are available from and/or extensively described in the *Handbook of Fluorescent Probes and Research Products* 8<sup>th</sup> Ed. (2001), available from Molecular Probes, Eugene, OR., as well as many other manufacturers.

Further amplification of the signal can be achieved by using combinations of specific binding members, such as antibodies and anti-antibodies, where the anti-antibodies bind to a conserved region of the target antibody probe, particularly where the antibodies are from different species. Alternatively specific binding ligand-receptor pairs, such as biotin-streptavidin, may be used, where the primary antibody is conjugated to one member of the pair and the other member is labeled with a detectable probe. Thus, one effectively builds a sandwich of binding members, where the first binding member binds to the cellular component and serves to provide for secondary binding, where the secondary binding member may or may not include a label, which may further provide for tertiary binding where the tertiary binding member will provide a label.

The secondary antibody, avidin, strepavidin or biotin are each independently labeled with a detectable moiety, which can be an enzyme directing a colorimetric reaction of a substrate having a substantially non-soluble color reaction product, a fluorescent dye (stain), a luminescent dye or a non-fluorescent dye. Examples concerning each of these options are listed below.

In principle, any enzyme that (i) can be conjugated to or bind indirectly to (e.g., via conjugated avidin, strepavidin, biotin, secondary antibody) a primary antibody, and (ii) uses a soluble substrate to provide an insoluble product (precipitate) could be used.

The enzyme employed can be, for example, alkaline phosphatase, horseradish peroxidase, beta-galactosidase and/or glucose oxidase; and the substrate can respectively be an alkaline phosphatase, horseradish peroxidase, beta.-galactosidase or glucose oxidase substrate.

Alkaline phosphatase (AP) substrates include, but are not limited to, AP-Blue substrate (blue precipitate, Zymed catalog p. 61); AP-Orange substrate (orange, precipitate, Zymed), AP-Red substrate (red, red precipitate, Zymed), 5-bromo, 4-chloro, 3-indolylphosphate (BCIP substrate, turquoise precipitate), 5-bromo, 4-chloro, 3-indolyl phosphate/nitroblue tetrazolium/iodonitrotetrazolium (BCIP/INT substrate, yellow-brown precipitate, Biomeda), 5-bromo, 4-chloro, 3-indolylphosphate/nitroblue tetrazolium (BCIP/NBT substrate, blue/purple), 5-bromo, 4-chloro, 3-indolyl phosphate/nitroblue tetrazolium/iodonitrotetrazolium (BCIP/NBT/INT, brown precipitate, DAKO, Fast Red (Red), Magenta-phos (magenta), Naphthol AS-BI-phosphate (NABP)/Fast Red TR (Red), Naphthol AS-BI-phosphate (NABP)/New Fuchsin (Red), Naphthol AS-MX-phosphate (NAMP)/New Fuchsin (Red), New Fuchsin AP substrate (red), p-Nitrophenyl phosphate (PNPP, Yellow, water soluble), VECTOR<sup>TM</sup> Black (black), VECTOR<sup>TM</sup>. Blue (blue), VECTOR<sup>TM</sup>. Red (red), Vega Red (raspberry red color).

Horseradish Peroxidase (HRP, sometimes abbreviated PO) substrates include, but are not limited to, 2,2' Azino-di-3-ethylbenz-thiazoline sulfonate (ABTS, green, water soluble), aminoethyl carbazole, 3-amino, 9-ethylcarbazole AEC (3A9EC, red). Alpha-naphthol pyronin (red), 4-chloro-1-naphthol (4C1N, blue, blue-black), 3,3'-diaminobenzidine tetrahydrochloride (DAB, brown), ortho-dianisidine (green), o-phenylene diamine (OPD, brown, water soluble), TACS Blue (blue), TACS Red (red), 3,3',5,5'-Tetramethylbenzidine (TMB, green or green/blue), TRUE BLUE<sup>TM</sup>. (blue), VECTOR<sup>TM</sup>. VIP (purple), VECTOR<sup>TM</sup>. SG (smoky blue-gray), and Zymed Blue HRP substrate (vivid blue).

Glucose oxidase (GO) substrates, include, but are not limited to, nitroblue tetrazolium (NBT, purple precipitate), tetranitroblue tetrazolium (TNBT, black precipitate), 2-(4-iodophenyl)-5-(4-nitrophenyl)-3-phenyltetrazolium chloride (INT, red or orange precipitate), Tetrazolium blue (blue), Nitrotetrazolium violet (violet), and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, purple). All tetrazolium substrates require glucose as a co-substrate. The glucose gets oxidized and the tetrazolium salt gets reduced and forms an insoluble formazan which forms the color precipitate.

Beta-galactosidase substrates, include, but are not limited to, 5-bromo-4-chloro-3-indoyl beta-D-galactopyranoside (X-gal, blue precipitate). The precipitates associated with each of the substrates listed have unique detectable spectral signatures (components).

5 The enzyme can also be directed at catalyzing a luminescence reaction of a substrate, such as, but not limited to, luciferase and aequorin, having a substantially non-soluble reaction product capable of luminescing or of directing a second reaction of a second substrate, such as but not limited to, luciferine and ATP or coelenterazine and Ca.sup.++, having a luminescing product.

10 The following references, which are incorporated herein provide additional examples. J.M Elias (1990) *Immunohistopathology: A practical approach to diagnosis*. ASCP Press (American Society of Clinical Pathologists), Chicago; J. F. McGinty, F. E. Bloom (1983) *Double immunostaining reveals distinctions among opioid peptidergic neurons in the medial basal hypothalamus*. Brain Res. 278: 145-153; and T. Jowett (1997) *Tissue In situ Hybridization: Methods in Animal Development*. John Wiley & Sons, Inc., New York; J Histochem Cytochem 1997 December 45(12):1629-1641.

15 Cellular preparations may be subjected to *in-situ* hybridization (ISH). In general, a nucleic acid sequence probe is synthesized and labeled with either a fluorescent probe or one member of a ligand:receptor pair, such as biotin/avidin, labeled with a detectable moiety. Exemplary probes and moieties are described in the preceding section. The sequence probe is complementary to a target nucleotide sequence in the cell. Each cell or cellular compartment containing the target nucleotide sequence may bind the labeled probe. Probes used in the analysis may be either DNA or RNA oligonucleotides or polynucleotides and may contain not only naturally occurring nucleotides but their analogs such as dioxynin dCTP, biotin dCTP 7-azaguanosine, azidothymidine, inosine, or uridine. Other useful probes include peptide probes and analogues thereof, branched gene DNA, peptidomimetics, peptide nucleic acids, and/or antibodies. Probes should have sufficient complementarity to the target nucleic acid sequence of interest so that stable and specific binding occurs between the target nucleic acid sequence and the probe. The degree of homology required for stable hybridization varies with the stringency

20  
25  
30

of the hybridization. Conventional methodologies for ISH, hybridization and probe selection are described in Leitch, et al. *In Situ Hybridization: a practical guide*, Oxford BIOS Scientific Publishers, Microscopy Handbooks v. 27 (1994); and Sambrook, J., Fritsch, E.F., Maniatis, T., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989).

5

The present invention is further illustrated by the following examples, which should not be construed as limiting in any way. The contents of all cited references are hereby expressly incorporated by reference.

#### **Example 1. Construction of Tissue Microarrays for a Survival Analysis of The Estrogen Receptor (ER) and HER2/neu and for Analysis of Nuclear Associated Beta-catenin**

Tissue microarray design: Paraffin-embedded formalin-fixed specimens from 345 cases of node-positive invasive breast carcinoma were identified. Areas of invasive carcinoma, away from *in situ* lesions and normal epithelium, were identified and three 0.6cm punch "biopsy" cores were taken from separate areas. Each core was arrayed into a separate recipient block, and five-micron thick sections were cut and processed as previously described (Konenen, J. et al., Tissue microarrays for high-throughput molecular profiling of tumor specimens, (1987) *Nat. Med.* 4:844-7). Similarly, 310 cases of colon carcinoma were obtained and arrayed, as previously described (Chung, G.G. et al., *Clin. Cancer Res.* (In Press)).

20

Immunohistochemistry: Pre-cut paraffin-coated tissue microarray slides were deparaffinized and antigen-retrieved by pressure-cooking (Kato, A.K. et al., (1997) *Biotech Histochem.* F2:291-8). Slides were stained with antibodies to one of three target antigens: monoclonal anti-E.R. (mouse, Dako Corporation, Carpinteria, CA), polyclonal anti-HER2/neu (rabbit, Dako Corp.), monoclonal (mouse clone 14, BD Transduction Labs, San Diego CA) anti-beta-catenin, or polyclonal rabbit anti-beta-catenin. Primaries were incubated overnight at 4°C. A corresponding goat antimouse or anti-rabbit secondary antibody conjugated to a horseradish peroxidase decorated dextran-polymer backbone was then applied for 1 hr (Envision, DAKO Corp.). Target antigens were either visualized with a visible light chromagen (Diaminobenzidine, DAKO) for visual analysis, or a fluorescent chromagen (Cy-5-tyramide, NEN Life Science Products, Boston, MA). Slides designated for automated analysis were

25

30

counterstained with DAPI for visualization of nuclei, and either polyclonal rabbit anticytokeratin (Zymed, So. San Francisco, CA) or rabbit anti-alpha-catenin (?) to distinguish between tumor cells and stroma as well as to visualize the cell membrane. In many cases, exponentially subtracted images of histospots stained with anti-cytokeratin provided an acceptable marker for the cell membrane due to the sub-membranous coalescence of cytokeratin in tumor cells. These antibodies were visualized using either Cy3- or Alexa 488-conjugated goat anti-mouse or anti-rabbit secondary antibodies (Amersham, Piscataway, NJ and Molecular Probes, Eugene, OR). Slides designated for visual inspection were counterstained with ammonium hydroxide acidified hematoxylin. Manual examination of microarrays for E.R., HER2/neu, and beta-catenin levels has been previously described (Snead, D.R. et al.,(1993) *Histopathology* 23:233-8).

Image analysis: Images of microarrays were obtained using a Deltavision platform and software (SoftWorx 2.5) (Applied Precision, Issaquah, WA), with an attached water-cooled Photometrics series 300 camera through a 10x Nikon Super-Fluor lens on a TE200 inverted fluorescent microscope with automated X,Y,Z stage movement. Low power images of microarrays were stitched together using multiple (~1500) low resolution images of the microarray (64x64 pixel). These images were analyzed by software algorithms described herein to determine the location of each . Subsequently, monochromatic, high resolution (1024x1024 pixel) images were obtained of each , both in the plane of focus and 8 microns below it. Image pairs for each fluorescent dye were obtained. Images were analyzed using additional algorithms as follows, in brief. Regions of interest (tumor) were identified using a mask derived from a ubiquitously-expressed epithelial-specific antigen (either cytokeratin or alpha-catenin). Images of fluorescently-tagged membrane and nuclear compartments were exponentially subtracted until a set amount of image intensity remained. Images were then combined so that there was minimal overlap of signal from one compartment to the next. Pixels in which a significant degree of overlap was present were negated from further analysis. The pixel intensity of exponentially subtracted images of the target antigen were assigned to one of three compartments: nuclear, membrane, or non-nuclear non-membrane (cytoplasm). Target intensities were analyzed as described below. For E.R. only nuclear-localized signal was used, for HER2/neu only membrane-localized signal was analyzed. For beta-catenin total signal, the ratio of nuclear to membrane signal, and the ratio of nuclear to total signal was analyzed.

Data analysis: staining scores from the breast cancers represent the averaged (for ER) or maximized (for HER2/neu) results from two scorable histospots. Subsequent studies revealed that analysis of a single histospot could provide significant statistical power to judge outcomes, so that staining scores from the colon cancer array represent the result of only one histospot. Overall survival analysis was assessed using Kaplan-Meier analysis and the Mantel-Cox log rank score for assessing statistical significance. Relative risk was assessed using the univariate Coxproportional hazards model. Analyses were performed using Statview 5.0.1 (SAS Institute, Cary NC).

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215